



# Coral Reefs and Climate Change

Denis Allemand  
(Centre Scientifique  
de Monaco)

Coral reefs are found in only a small percentage of global oceans, between 0.08 and 0.16%, but they shelter about one third of the marine species known today. This ecological success is due to a symbiosis between a coral and an intracellular microalgae, commonly called zooxanthellae. "Organismic engineers", they are the source of the largest biological constructions on the planet. Genuine oases of life, they support the direct sustenance of more than 500 million people in the world from fishing, but they engage human interest also for other reasons: protection of coasts against erosion, high value tourist areas... Ecological services from coral reefs are estimated at approximately 30 billion USD per year. Their growth is dependent on many factors (light, temperature, pH, nutrients, turbidity...). They are therefore extremely sensitive to the current changes in our environment: water temperature variability, ocean acidification, in addition to localized disruptions (pollution, sedimentation, coastal development, overfishing, marine shipping...). An increase of less than 1 degree above a threshold value is sufficient to cause bleaching. It breaks the coral symbiosis with their zooxanthellae throughout the populations, leading to the disappearance of the reef. Similarly, ocean acidification disrupts the formation of a coral's skeleton, and many other biological functions such as reproduction. We actually estimate that approximately 20% of the global coral reefs have already disappeared completely; 25% are in high danger; and 25% more will be threatened by 2050 if positive management action is not taken.

## WHAT IS A CORAL REEF?

Coral reefs are ecosystems typically found in shallow waters of the intertropical zone (approximately between 33° North and 30° South). The three-dimensional architecture of this ecosystem is formed by the buildup of calcareous skeletons of marine organisms called reef-building corals (Cnidaria, Scleractinia). They are cemented together by the biological activity of calcareous organisms (macroalgae, sponges, worms, molluscs...). Corals are named "engineering organisms", while the reef is considered "biogenic" because they are the result of biological activity. Coral reefs therefore represent ecosystems that have been built by their own inhabitants.

The total area covered by coral reefs varies, depending on the calculation methods, between 284,300km<sup>2</sup> (Smith, 1978) and 617,000km<sup>2</sup> (Spalding *et al.*, 2001), therefore covering between 0.08 and 0.16% of the surface of the ocean. French reefs alone cover an area of 55,557km<sup>2</sup>. The largest reef is the Great Barrier Reef which runs along the north-eastern coast of Northern Australia over a distance of 2300 km. It is reputed to be the only animal construction visible from space. The second largest reef is French, the New Caledonian barrier, which is 1600 km long. These two barrier reefs have been registered by the UNESCO World Heritage (respectively in 1981 and 2008).



Coral reefs come in different shapes and sizes, the first published description dating from Charles Darwin during his voyage on the Beagle (Darwin, 1842):

- Fringing reefs: They follow the coastline, maintaining an active growth area offshore and an accumulation of dead coral inshore, forming a platform reef that over time turns into a lagoon.
- Barrier reefs: the fringing reef becomes a barrier reef subsequent to the progressive sinking of an island. In this way, the lagoon becomes larger and the reef can reach up to 1km away from the coast.
- Atolls: these are the ultimate step in the evolution of a reef, where the island has completely disappeared below the sea surface. Atolls preserve the initial circular shape of the island. There are approximately 400 atolls in the world.

Reef growth is of the order of about 4kg of calcium carbonate ( $\text{CaCO}_3$ ) per  $\text{m}^2$  per year (Smith & Kinsey, 1976), but values can vary considerably from one reef to another, in some cases reaching up to 35kg  $\text{CaCO}_3/\text{m}^2/\text{year}$  (Barnes & Chalker, 1990), *i.e.* a vertical annual growth rate of 1 to 7mm. Many factors influence these growth rates: light, temperature (optimal between 22° and 29°C), nutrients, currents, turbidity, pH and the saturation state of calcium carbonate in the seawater...

The formation of calcium carbonate by reef-building organisms causes the release of carbon dioxide into the surrounding environment. Hence, contrary to what has been believed, a reef mainly dominated by coral acts as a minor source and not as a sink of  $\text{CO}_2$  (about 1.5mmol  $\text{CO}_2/\text{m}^2$  day. Tambutté *et al.*, 2011 for a review). Nevertheless, reefs still do play an important role as a carbon sink (as  $\text{CaCO}_3$ ), with rates of the order of 70 to 90 million tonnes of carbon per year (Frankignoulle & Gattuso, 1993).

## CORALS, AT THE ORIGIN OF THE REEF

Reefs are mainly built by corals. Formerly known as stony corals, reef-building corals are now in-

cluded in the Order of Scleractinians (subclass *Hexacorallia*, class *Anthozoa* of phylum *Cnidaria*). Among the *Scleractinia*, about half the amount of species (about 660 out of 1,482 species known to date, Cairns, 1999) are involved in reef construction. These are called hermatypic. They consist of polyps of variable sizes, depending on the species, and form functional units. Each polyp has a mouth surrounded by tentacles. The polyps are connected to each other by network of cavities, the coelenteron, which covers the coral tissue. The whole assemblage is known as colonial (even though the colony functions as a single organism) while individual corals are called modular animals. They present various shapes and sizes, depending on whether the species are branching coral, blade coral, encrusting, or massive coral for example, and show growth rates that can exceed 15cm per year of axial growth in their natural environment (Dullo, 2005). The size of certain massive corals may even exceed 6m in diameter.

The degree of success for a reef to develop and to thrive is mainly related to the capability of the majority of scleractinian corals (just under 900 species, Michel Pichon, Comm. Pers.) to establish a mutual symbiosis with photosynthetic dinoflagellates commonly called *zooxanthellae* (*e.g.* *Symbiodinium sp.*). These microalgae reside inside the coral's gastroderm, isolated from the animal's cytoplasm by a perisymbiotic membrane that regulates the exchanges between the symbionts and the host (Furla *et al.*, 2011 for a review). These two partners have co-evolved since the Triassic (Muscattine *et al.*, 2005), developing unique abilities (*e.g.* the ability for the hosts to actively absorb  $\text{CO}_2$  and nutrients and to protect themselves from ultraviolet rays, hyperoxia and oxidative stress; the ability of the algal symbiont to exchange nutrients with its host; Furla *et al.*, 2005, 2011). Due to the presence of zooxanthellae, the distribution of corals at depth is dependent upon light availability (generally between 0 and 30m depth). Thanks to modern sequencing techniques, a strong diversity in bacteria has been evidenced inside corals. These bacteria appear to play an important physiological role. The entire community of these living organisms forms a functional unit called a holobiont, often referred to as a super-organism.



Symbiont photosynthesis is also related to another function of coral, biomineralization, that is to say its ability to build a limestone or biomineral skeleton. The property of a biomineral is that it is a composite material, comprising both a mineral fraction and an organic fraction. Even though the latter is minimal (<1% by weight), it plays a key role in controlling the deposition of calcium carbonate in the form of aragonite (German *et al.*, 2011, Tambutté *et al.*, 2008, 2011). Using mechanisms that are still a matter of debate, light, via symbiont photosynthesis, has been observed to stimulate the calcification of coral by a factor reaching 127 in comparison to night calcification. However, in most cases, this factor varies between 1 and 5, with an average value of 4 (Gattuso *et al.*, 1999).

Coral reproduction is typically sexual and involves a larval stage called *planula* which ensures the species dispersion. They can also have a high asexual reproductive capacity by fragmentation. This capacity is utilized in the development of *ex situ* cultures.

## CORAL AND CORALS

Coral, in its name, hides many organisms belonging to the *Cnidaria* phylum and is at the base of particular ecosystems:

- Cold-water corals, also called deep-sea corals: these corals belong to the same order of cnidarians as reef-building corals. Like them, they are engineering organisms, capable of building a rich ecosystem that provides habitat for many other creatures in the deep waters of the Atlantic, Pacific, as well as the Mediterranean Sea. Unlike their surface water cousins, they are acclimated to cold waters (6° -14°C) and do not host photosynthetic algae. These reefs therefore play a significant role as shelters and nursery areas for many species of fish of commercial interest (Roberts *et al.*, 2009).
- The coralligenous in the Mediterranean: they are formed by an assemblage of stationary creatures (e.g. gorgonians, red coral, encrusting calcareous algae...). The co-

ralligenous in the Mediterranean form a very rich coastal ecosystem, especially along underwater cliffs. It is of particular interest both for fishing and aquatic tourism (RAC/SPA 2003).

## THE CORAL REEF: A BIODIVERSITY HOT-SPOT

The ability to live in symbiosis with dinoflagellates has allowed coral reefs to build large constructions in usually oligotrophic conditions, that is to say, nutrient-poor waters. Coral reefs have existed since the Triassic, about 200 million years ago. However, since that time there have been many phases of disappearance/reappearance. The development of the Great Barrier Reef seems to have begun 20 million years ago. However, primitive forms that are different from modern corals, have existed long before the Triassic, during the Devonian about 400 million years ago. Coral reefs are home to the greatest biodiversity on Earth with 32 of the 34 animal phyla known to date and include a third of marine species known so far, representing nearly 100,000 species (Porter & Tougas, 2001). Hence, 30% of the known marine biodiversity is sheltered in less than 0.2% of the total surface of the oceans! In the marine environment, they therefore represent the equivalent of the primary tropical forests. For comparison, the number of species of molluscs found on 10 m<sup>2</sup> of reef in the South Pacific is greater than what has been acknowledged throughout the whole North Sea. As another example, in New Caledonia there are over 400 species of coastal nudibranchs while in mainland France there is little more than a dozen species for an equivalent coastline.

This "biodiversity" is however not homogeneous between reefs. Indeed, there is a skewed distribution of the diversity and abundance of corals between the Atlantic and Pacific Oceans, as well as within these oceans. In these two oceans, the diversity and abundance are concentrated in the western parts: the Coral Triangle (also called "Centre for Coral Biodiversity") in the Pacific, including the -Indonesia Malaysia - Philippines - China Sea - Solomon Islands region; the



Caribbean in the Atlantic. There is also a strong east-west longitudinal gradient. The fauna and flora associated with reefs generally follow similar gradients.

## THE CORAL REEF: AN EXCEPTIONAL WEALTH FOR MANKIND

Coral reefs border the coasts of more than 80 countries across the world (Sheppard *et al.*, 2009) for which they represent an important source of income, just as much in terms of a food resources, coastal protection and tourism... Approximately 275 million people worldwide live within 30km of a coral reef and the livelihood of over 500 million people directly depend on reefs. On one hand economists estimate that the annual value of the benefits provided by the reefs is worth slightly more than 24 billion euros (Chen *et al.*, 2015). On another hand, the TEEB report (TEEB, 2010) has estimated that the destruction of coral reefs would represent a loss of about € 140 billion per year.

The ecosystemic benefits provided by coral reefs include:

### 1. Natural resources

- Food: coral reefs provide 9 to 12% of the world catch of edible fish and 20 to 25% of the fish catch in developing countries (Moberg & Folke, 1999). This figure reaches 70 to 90% for the South East Asian countries (Garcia & de Leiva Moreno, 2003). The total estimated income of reef fisheries is about 5 billion euros (Conservation International, 2008). Most of these fisheries are traditional, carried out on foot by the local population, especially women and children who collect fish, molluscs (clams), crustaceans (crabs and lobsters) and sea cucumber (also referred to as trepang). A healthy reef is estimated to annually provide 5 to 10 tonnes of fish and invertebrates per km<sup>2</sup>.
- Mineral resources: coral reefs provide housing construction materials (Maldives, Indonesia), sand for the construction of roads or fertilizers for agricultural land. Coral reefs in the Maldives thus supply about 20,000m<sup>3</sup>

of material annually (Moberg & Folke, 1999).

- Live Resources: beyond fishing for food needs, reefs also represent a fishing reserve for coral reef aquariology (15 million fish per year for 2 million aquarists in the world) and pearl farming, etc.

### 2. Conservation

- Coastal Protection: coral reefs have an undeniable role in the protection of coastline from the destructive action of waves and tsunamis. More than 150,000 km of coastline are naturally protected by barrier reefs (<http://www.coralguardian.org>). A typical coral reef can absorb up to 90% of the impact load of a wave (Wells, 2006). During the devastating 2004 tsunami in the Indian Ocean, coasts protected by healthy coral reefs were much less affected by the deadly wave. The value of coastal protection against natural disasters has been estimated to lie between 20,000 and 27,000 euros per year per hectare of coral (TEEB, 2010). The total profit is estimated at 7 billion euros per year (Conservation International, 2008).

### 3. Cultural resources

- Tourism: tourists are attracted to the natural beauty of coral reefs (via terrestrial tourism, diving). The large number of visitors promotes employment, a windfall for the poverty-stricken parts of the world. For example, the Australian Great Barrier Reef attracts about 2 million visitors annually, producing an income of around 4 billion Euros for the Australian economy and 54,000 jobs (Biggs, 2011). According to estimates compiled by the TEEB report, one hectare of coral reef represents a yearly profit of 64,000 to 80,000 Euros from tourism and recreational opportunities. Eco-tourism alone earned 800,000 euros per year in the Caribbean. The total annual income from coral reefs is estimated around 8 billion euros (Conservation International, 2008).
- Cultural or religious heritage: Coral reefs are at the base of many cultural and religious traditions. In southern Kenya, for example, many religious rituals are structured around coral reefs in order to appease the spirits (Moberg & Folke, 1999).



- Medical resources: the numerous marine invertebrates (sponges, molluscs or soft corals) represent a potential supply of new drugs for human health. Coral is also starting to be used as a biological model to better understand immunity or aging mechanisms (Moberg & Folke, 1999).

## THE CORAL REEF: LOCAL AND GLOBAL THREATS

The coral reef ecosystems are currently threatened both locally (pollution, sedimentation, unsustainable coastal development, nutrient enrichment, overfishing, use of destructive fishing methods...) and, since the 1980s, globally (global warming, ocean acidification). The Global Coral Reef Monitoring Network (GCRMN) estimates that at present, 19% of reefs have been destroyed, 15% are seriously damaged and may disappear within the next ten years, and 20% could disappear within less than 40 years. More positively, 46% of the world's reefs are still healthy (Wilkinson, 2008). The rare monitoring studies on reef growth show a clear long-term decrease in coral cover: in an analysis of 2258 measurements from 214 reefs of the Great Barrier during the 1985-2012 period, De'ath *et al.*, (2012) evidenced a decline in the coral cover from 28.0% to 13.8% as well as loss of 50.7% of initial coral cover.

Among the global events that affect coral reefs, the increasing temperature of surface water is causing a widespread phenomenon, coral bleaching. Unique example, visible to the naked eye, of the impact of climate change on an ecosystem, coral bleaching is the result of the rupture of the symbiosis between corals and zooxanthellae symbionts. Although it can be reversible during the first few days, this bleaching effect inevitably leads to coral death a few weeks after the symbiosis is halted (Hoegh-Guldberg, 1999; Weis &

Allemand, 2009). This phenomenon, whose inner mechanisms are still under debate, usually occurs when the temperature exceeds a certain threshold by 0.5°C.

A second event is just as seriously affecting coral biology: ocean acidification, also referred to as the other effect of CO<sub>2</sub> (Doney *et al.*, 2009). Part of the excess carbon dioxide produced by human activities dissolves into the oceans, reducing on one hand the greenhouse effect (and thus reducing the increase in global temperature), but on the other hand causing an increasing acidity of the oceans, according to the following reaction:



To date, the pH of seawater has decreased by about 0.1 units since the beginning of last century (from 8.2 to 8.1) which corresponds to an increase in the acidity of the water by about 30% (Gattuso & Hansson, 2011). Acidification primarily affects the calcification rates of corals, and therefore reef growth. However, it appears that the effects vary greatly from one species to another (Erez *et al.*, 2011). The differences in sensitivity may be due to a differential ability of the animal to control the pH of its calcification site (Holcomb *et al.*, 2014; Venn *et al.*, 2013). However the increase in dissolved CO<sub>2</sub> has also been found to cause many other effects on coral physiology, including the alteration of gene expression (Moya *et al.*, 2012; Vidal-Dupiol *et al.*, 2013).

Unfortunately, our present knowledge of the physiology of these creatures is too insufficient to predict whether corals will be able to adapt to rapid changes in the environment, especially since earlier studies suggest that the combined effects of the decrease in the pH with the increase in temperature of the sea seem to have cumulative effects (Reynaud *et al.*, 2003).





## REFERENCES

- ALLEMAND D., FURLA P. and BÉNAZET-TAMBUTTÉ S., 1998 – *Mechanisms of Carbon Acquisition for Endosymbiont Photosynthesis in Anthozoa*. Can J Bot 76: 925-941.
- ALLEMAND D., TAMBUTTÉ É., ZOCCOLA D. and TAMBUTTÉ S., 2011 – *Coral Calcification, Cells to Reefs*. In *Coral Reefs: an Ecosystem in Transition*. Springer Netherlands.
- BARNES D. J. and CHALKER B. E., 1990 – *Calcification and Photosynthesis in Reef-Building Corals and Algae*. In *Coral Reefs*. Amsterdam: Elsevier.
- BIGGS D., 2011 – *Understanding Resilience in a Vulnerable Industry: the Case of Reef Tourism in Australia*. Ecology and Society 16 (1): 30.
- CAIRNS S. D., 1999 – *Species Richness of Recent Scleractinia*. Atoll Res Bull 459: 1-46.
- CAR/ASP, 2003 – *Le coralligène en Méditerranée*. PNUE.
- CHEN P. Y., CHEN C. C., CHU L. and MCCARL B., 2015 – *Evaluating the Economic Damage of Climate Change on Global Coral Reefs*. Global Environmental Change 30: 15-20.
- CONSERVATION INTERNATIONAL, 2008 – *Economic Values of Coral Reefs, Mangroves, and Seagrasses: a Global Compilation*. Center for Applied Biodiversity Science, Arlington.
- DARWIN C. R., 1842 – *The Structure and Distribution of Coral Reefs. Being the First Part of the Geology of the Voyage of the Beagle, under the Command of Capt. Fitzroy, R.N. during the Years 1832 to 1836*. London: Smith Elder and Co.
- DE'ATH G., FABRICIUS K. E., SWEATMAN H. and PUOTINEN M., 2012 – *The 27-Year Decline of Coral Cover on the Great Barrier Reef and its Causes*. Proceedings of the National Academy of Sciences of the United States of America, 109 (44), 17995-17999.
- DONEY S. C., V. FABRY J., FEELY R. A. and KLEYPAS J. A., 2009 – *Ocean Acidification: the Other CO<sub>2</sub> Problem*. Ann Rev Marine Sci 1: 169-192.
- DULLO W. C., 2005 – *Coral Growth and Reef Growth: a Brief Review*. Facies 51: 33-48.
- EREZ J., REYNAUD S., SILVERMAN J., SCHNEIDER K. and ALLEMAND D., 2011 – *Coral Calcification under Ocean Acidification and Global Change*. In *Coral Reefs: an Ecosystem in Transition*. Springer Netherlands.
- FRANKIGNOULLE M. and GATTUSO J.-P., 1993 – *Air-Sea CO<sub>2</sub> Exchange in Coastal Ecosystems*. NATO ASI Series 14: 233-248.
- FURLA P., ALLEMAND D., SHICK M., FERRIER-PAGÈS C., RICHIER S. et al., 2005 – *The Symbiotic Anthozoan: a Physiological Chimera between Alga and Animal*. Integr Comp Biol 45: 595-604.
- FURLA P., RICHIER S. and ALLEMAND D., 2011 – *Physiological Adaptation to Symbiosis in Cnidarians*. *Coral Reefs: an Ecosystem in Transition*. Springer Netherlands.
- GARCIA S. M. and DE LEIVA MORENO J. I., 2003 – *Global Overview of Marine Fisheries*. In *Responsible Fisheries in the Marine Ecosystem*. FAO & CABI Publishing.
- GATTUSO J.-P., ALLEMAND D. and FRANKIGNOULLE M., 1999 – *Photosynthesis and Calcification at Cellular, Organismal and Community Levels*. In *Coral Reefs: a Review on Interactions and Control by Carbonate Chemistry*. Am Zool 39: 160-183.
- GATTUSO J.-P., FRANKIGNOULLE M. and WOLLAST R., 1998 – *Carbon and Carbonate Metabolism in Coastal Aquatic Ecosystems*. Annu Rev Ecol Syst 29: 405-433.
- HOEGH-GULDBERG O., 1999 – *Climate Change, Coral Bleaching and the Future of the World's Coral Reefs*. Mar Freshwater Res 50: 839-866.
- HOLCOMB M., VENN A. A., TAMBUTTÉ É., TAMBUTTÉ S., ALLEMAND D. et al., 2014 – *Coral Calcifying Fluid Ph Dictates Response to Ocean Acidification*. Sci Rep 4: 5207.
- HOULBRÈQUE F. and FERRIER-PAGES C., 2009 – *Heterotrophy in Tropical Scleractinian Corals*. Biol Rev. 84: 1-17.
- MOBERG F. and FOLKE C., 1999 – *Ecological Goods and Services of Coral Reef Ecosystems*. Ecol Econ 29: 215-233.
- MOYA A., HUISMAN L., BALL E. E., HAYWARD D. C., GRASSO L. C. et al., 2012 – *Whole Transcriptome Analysis of the Coral Acropora millepora Reveals Complex Responses to CO<sub>2</sub>-driven Acidification during the Initiation of Calcification*. Mol Ecol 21: 2440-2454.



- MUSCATINE L., GOIRAN C., LAND L., JAUBERT J., CUIF J. P. *et al.*, 2005 – *Stable Isotopes ( $^{13}\text{C}$  and  $^{15}\text{N}$ ) of Organic Matrix from Coral Skeleton*. Proc Natl Acad Sci USA 102: 1525-1530.
- PORTER J. W. and TOUGAS J. I., 2001 – *Reef Ecosystems: Threats to their Biodiversity*. In *Encyclopedia of Biodiversity*. San Diego: Academic Press.
- REYNAUD S., LECLERCQ N., ROMAINE-LIOUD S., FERRIER-PAGÈS C., JAUBERT J. *et al.*, 2003 – *Interacting Effects of  $\text{CO}_2$  Partial Pressure and Temperature on Photosynthesis and Calcification in a Scleractinian Coral*. Global Change Biol 9: 1660-1668.
- TAMBUTTÉ S., HOLCOMB M., FERRIER-PAGÈS C., REYNAUD S., TAMBUTTÉ É. *et al.*, 2011 – *Coral Biomineralization: from the Gene to the Environment*. J Exp Mar Biol Ecol: 58-78, 2011.
- SMITH S. V. and KINSEY D. W., 1976 – *Calcium Carbonate Production, Coral Reef Growth, and Sea Level Change*. Science 194: 937-939.
- TAMBUTTÉ S., TAMBUTTÉ É., ZOCCOLA D. and ALLEMAND D., 2008 – *Organic Matrix and Biomineralization of Scleractinian Corals*. In *Handbook of Biomineralization*. Wiley-VCH Verlag GmbH.
- TEEB, 2010– *The Economics of Ecosystems and Biodiversity Ecological and Economic Foundations*. Pushpam Kumar, Earthscan.
- VENN A. A., TAMBUTTÉ É., HOLCOMB M., LAURENT J., ALLEMAND D. *et al.*, 2013 – *Impact of Seawater Acidification on Ph at the Tissue-Skeleton Interface and Calcification in Reef Corals*. Proc Natl Acad Sci USA 110: 1634-1639.
- VIDAL-DUPIOL J., ZOCCOLA D., TAMBUTTÉ É., GRUNAU C., COSSEAU C. *et al.*, 2013 – *Genes Related to Ion-Transport and Energy Production Are Upregulated in Response to  $\text{CO}_2$ -Driven Ph Decrease in Corals: New Insights from Transcriptome Analysis*. PLoS One 8: e58652.
- WEIS V. M. and ALLEMAND D., 2009 – *What Determines Coral Health?* Science 324: 1153-1155.
- WELLS S., 2006 – *In The Front Line Shoreline Protection and other Ecosystem Services from Mangroves and Coral Reefs*. UNEP-WCMC Biodiversity Series 24: 1-34.
- WELLS S., 2006 – *Shoreline Protection and other Ecosystem Services from Mangroves and Coral Reefs*. UNEP-WCMC Biodiversity Series 24.